

Evaluation of Bacterial Species to Determine Antimicrobial Resistance in Patients with Chronic Rhinosinusitis after Surgery of Paranasal Sinuses Referring to Amiralmomenin Hospital in Rasht, 2018

Shadman Nemati¹, Ali Mojtahedi², Soheil Soltanipour³, Masoumeh Sharifigar Mavari⁴, Samaneh Rouhi⁵

1. Professor, Specialty of Otolaryngology and Head and Neck Surgery, Otorhinolaryngology Research Center, Department of Otolaryngology and Head and Neck Surgery, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran. ORCID ID: 0000-0001-8536-4703

2. Associate Professor of Medical Microbiology, Cellular and Molecular Research Center, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran. ORCID ID: 0000-0002-5385-9367

3. Associate Professor of Community Medicine, Eye Research Center, Department of Eye, Amiralmomenin Hospital, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran. ORCID ID: 0000-0001-7768-1121

4. General Practitioner, Otorhinolaryngology Research Center, Department of Otolaryngology and Head and Neck Surgery, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran. ORCID ID: 0000-0003-0943-3710

5. PhD of Medical Bacteriology, Medical Microbiology Research Center, Qazvin University of Medical Sciences, Qazvin, Iran., (Corresponding Author), Tel: 028-833790622, Email: roohi.sanmaneh@yahoo.com, ORCID ID: 0000-0003-0160-0924

ABSTRACT

Background and Aim: Bacterial resistance to antibiotics has made treatment difficult. The purpose of this study was to investigate bacterial species in patients with chronic rhinosinusitis after surgery of paranasal sinuses to determine antimicrobial resistance patterns of them.

Materials and Methods: The data of 70 patients after paranasal sinuses surgery in Amiralmomenin hospital in Rasht city, in 2018 were evaluated. The identification of bacteria by microbiological laboratory methods and microbial susceptibility test was performed by disk diffusion method. For data analysis, SPSS version 22 software and chi-square test were used ($p \leq 0.05$).

Results: 62 (88.57%) positive bacterial culture samples were identified. The most abundant strains was *Staphylococcus epidermidis* (38.70%). *Staphylococcus aureus* had the highest antibiotic resistance to penicillin and oxacillin (52.94%) and *Staphylococcus epidermidis* to penicillin (62.50%). Highest antibiotic resistance in *Pseudomonas aeruginosa* was to ceftazidime (90.90%). *Escherichia coli* was resistant to ceftazidime and ampicillin (100%) and *Hafnia alvei* was resistant to ceftazidime (100%). *Klebsiella aerogenes* had higher resistant to ceftazidime and cefixime (100%). With increasing of patient's age, resistance to antibiotics increased ($p \leq 0.05$).

Conclusion: Antibiotic resistance was observed in bacterial samples isolated from patients after surgery. Given that antibiotic resistance may cause failure in the treatment. Monitoring of the antibiotic-resistant pattern is necessary to select the appropriate antibiotic.

Keywords: Bacterial Species, Antimicrobial Resistance Pattern, Chronic Rhinosinusitis, Paranasal Sinuses

Received: July 27, 2019

Accepted: Jan 7, 2020

How to cite the article: Shadman Nemati, Ali Mojtahedi, Soheil Soltanipour, Masoumeh Sharifigar Mavari, Samaneh Rouhi. Evaluation of Bacterial Species to Determine Antimicrobial Resistance in Patients with Chronic Rhinosinusitis after sSurgery of Paranasal Sinuses Referring to Amiralmomenin Hospital in Rasht, 2018. SJKU 2020; 25(2): 1-13.